



Evaluating Carbon Stocks (2020-2023) in *Rhizophora mucronata* Plantations under the PHE ONWJ REMAJA Program in Pantai Bahagia Villages, Bekasi Regency, Indonesia

Judul Jurnal

Rahman ^a, Luisa Febrina Amalo ^{b*}, Pungki Ari Wibowo ^b,
Luluk Dwi Wulan Handayani ^b, Hadi Supardi ^c,
L. M. Alfin Agushara Bena ^c and Arif Rahman ^c

^a Department of Marine Science, Faculty of Fisheries and Marine Science, Pattimura University, Indonesia.

^b The Central for Environment Research, IPB University, West Java, Bogor, Indonesia.

^c Department of Environmental, HSSE Division, PT. Pertamina Hulu Energi Offshore North West Java, Jakarta, Indonesia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajee/2024/v23i10615>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124263>

Original Research Article

Received: 14/07/2024

Accepted: 30/09/2024

Published: 03/10/2024

*Corresponding author: E-mail: luisafebrina@gmail.com;

Cite as: Rahman, Luisa Febrina Amalo, Pungki Ari Wibowo, Luluk Dwi Wulan Handayani, Hadi Supardi, L. M. Alfin Agushara Bena, and Arif Rahman. 2024. "Evaluating Carbon Stocks (2020-2023) in *Rhizophora Mucronata* Plantations under the PHE ONWJ REMAJA Program in Pantai Bahagia Villages, Bekasi Regency, Indonesia". *Asian Journal of Environment & Ecology* 23 (10):121-29. <https://doi.org/10.9734/ajee/2024/v23i10615>.

ABSTRACT

This study evaluates the carbon storage potential of *Rhizophora mucronata* plantations as part of the North Java Coast Mangrove Restoration Program (REMAJA) implemented by PT. Pertamina Hulu Energi Offshore North West Java (PHE ONWJ) in Pantai Bahagia Village, Bekasi Regency. A total of 8,000 mangrove seedlings were planted in 2020, and the diameter growth and carbon stock were monitored from 2020 to 2023. The monitoring process involved measuring the trunk diameter of the mangrove, combined with allometric equations to estimate biomass. Carbon sequestration was calculated by applying conversion factors to the biomass values, allowing the quantification of stored carbon. This method provides an accurate assessment of the mangrove's role in carbon sequestration over time. The results showed that the survival rate of *Rhizophora mucronata* seedlings was exceptionally high, averaging over 98%. The total carbon stock increased annually, reaching 15.59 tons of carbon (C) in 2023, 9.57 tons in AGC, and 6.02 tons in BGC. The result represents a significant increase compared to previous years, with an additional 5.14 tons of carbon compared to 2022, 13.75 tons more than in 2021, and 15.01 tons more than in 2020. Based on these results, the carbon dioxide sequestration contribution of *R. mucronata* during the periods of 2020-2021, 2021-2022, and 2022-2023 was 4.62 tons CO_{2e}, 31.58 tons CO_{2e}, and 18.83 tons CO_{2e}, respectively. In conclusion, the *R. mucronata* plantation under this program significantly enhanced carbon stocks, supporting climate change mitigation through long-term carbon storage. This program demonstrates that mangrove rehabilitation can play a crucial role in protecting coastal ecosystems and reducing greenhouse gas emissions.

Keywords: Carbon stock; climate change mitigation; *R. mucronata*; mangrove rehabilitation.

1. INTRODUCTION

Mangrove ecosystems are the main blue carbon ecosystems found in coastal areas. These ecosystems have varying blue carbon potentials influenced by the density, canopy cover, and diameter of mangroves [1]. The potential carbon stock of mangroves is 956 MgC ha⁻¹ [2] and even reaches 1082.55 MgC ha⁻¹ [3]. A recent study by Rahman et al. [1] found that the total potential carbon stock reaches 1619.88 MgC ha⁻¹. The potential of carbon stocks is much greater than those in other ecosystems, such as seagrasses, saline swamps, and tropical rainforests [4,5]. Based on this potential, blue carbon has become a new paradigm for sustainable mangrove management [6].

Indonesia's coastal area is one of the world's most significant mangrove ecosystem habitats. This area had a mangrove area of 3.12 million ha in 1996 and 2.95 million ha in 2020 [1]. The decrease in area is caused by the conversion of land into ponds, settlements, and timbers [7] which has an impact on the decrease in the density and carbon sequestration potential of several mangrove species, especially from the *Rhizophora* and *Sonnerartia* species [8].

Deforestation of mangrove ecosystems in Indonesia occurs in almost all coastal areas of Indonesia, especially the island of Java, which is

known as the Bekasi Regency. Amalo et al. [9] reported that the mangrove ecosystem in Bekasi Regency has decreased by 85.58 ha triggered by the conversion of land into ponds, so conservation efforts are needed to maintain existing mangroves, for example as ecotourism and potential land rehabilitation for an increase in area and density.

The success of mangrove rehabilitation efforts can be achieved when considering ecological factors, primarily related to the adaptability of mangroves to environmental conditions such as temperature, salinity, and waves [10]. One of the mangrove species that has high adaptability to the ecological environment is *Rhizophora mucronata* [11,12]. It is indicated by the existence of this species, commonly found in almost all coastal areas of Indonesia [13,14].

Based on this, PT. Pertamina Hulu Energy Offshore North West Java (PT. PHE ONWJ) carries out mangrove rehabilitation on the coast of Bekasi Regency, especially Pantai Bahagia Village through the North Java Coast Mangrove Restoration (REMAJA) program. A total of 8000 *Rhizophora mucronata* seedlings were planted in 2020. The planting aims to improve the quality of the mangrove ecosystem, including biodiversity, area, density, and ecosystem services, as well as the potential for carbon sequestration in climate change mitigation efforts. Therefore, this study

aims to evaluate the potential carbon storage from planting *Rhizophora mucronata* by monitoring diameter growth from 2020 to 2023.

strategy focused on optimizing survival rates by ensuring that the seedlings were well-adapted to the local substrate and hydrological conditions.

2. METHODS

2.1 Description of Rehabilitation Program

Mangrove rehabilitation was carried out in 2020 by PT. PHE ONWJ on the coast of Pantai Bahagia Village, Bekasi Regency (Fig. 1). A total of 8,000 *R. mucronata* seedlings were planted in 2020 across the potential rehabilitation area, characterized by sandy mud substrate and favorable environmental conditions. The seedlings selected for planting were carefully chosen based on their ecological suitability to the coastal area of Pantai Bahagia Village, where *R. mucronata* naturally occurs. The seedlings used in this rehabilitation effort were approximately 6–12 months old, with an average height of 30–50 cm. These attributes were considered ideal for promoting rapid establishment and growth in the given environmental conditions. The planting

2.2 Measurement of Mangrove Diameter

Mangrove diameter measurements were conducted annually from 2020 to 2023. Three hundred (300) mangrove seedlings, selected from 8,000 planted seedlings, were measured using a stratified random sampling method. The selection of 300 seedlings ensured the representation of the ecological characteristics across the planting area. The data obtained from these measurements included each year's average diameter and growth rate.

The initial diameter of *R. mucronata* seedlings was measured at the point where the propagule meets the stem, particularly for seedlings with a height of ≤ 50 cm [15]. For seedlings that have developed into saplings, the diameter was measured 50 cm above the ground [16].




Fig. 1. Map of the *R. mucronata* planting by PHE ONWJ

2.3 Data Analysis

2.3.1 Survival Rate

The survival rate of *R. mucronata* mangrove is the percentage of the ratio of live seedlings to the total planted seedlings. However, to represent 8000 seedlings, 300 were selected proportionally by paying attention to the representation of each ecological character of the mangrove planting area. Mathematically, the survival rate equation can be written as follows:

$$R (\%) = \frac{\sum SLt}{N} \times 100 \%$$

R is the survival rate (%); $\sum SLt$ is the number of seedlings alive during observation; and N is the total number of seedlings planted.

2.3.2 Mangrove Biomass

Evaluation of *R. mucronata* biomass was carried out on diameter growth in the first year (2020-2021), second (2021-2022), and third (2022-2023). Biomass was estimated based on above-ground biomass (AGB) and below-ground biomass (BGB). The allometric equation used for biomass estimation refers to Komiyama et al. [16] as follows:

$$\begin{aligned} \text{Above-ground biomass (kg)} &= 0.251\rho D^{2.46} \\ \text{Below-ground biomass (kg)} &= 0.199\rho^{0.899}D^{2.22} \end{aligned}$$

Notes: ρ is mangrove wood density, *R. mucronata* is 0.8483 based on data from the World Agroforestry Center [17].

2.3.3 Carbon stock and CO₂ absorption

Carbon stock estimation is carried out by multiplying the value of biomass by the carbon

fraction. The magnitude of the carbon fraction value refers to Rahman et al. [18], which is 0.4682. Meanwhile, the carbon absorption potential (CO₂e) is obtained by multiplying the potential carbon stock with the value of the comparison of the molecular mass of CO₂ (Mr. CO₂ = 44 grams/mol) to the atomic mass of C (Ar C = 12 grams/mol). Mathematically, the equation for stock estimation and carbon sequestration is as follows:

$$\begin{aligned} CS &= B \times 0,4682 \\ CO_2e &= CS \times (44/12) \end{aligned}$$

Information:

CS = Carbon Stock (kg/stand)
 B = biomass (kg/stand) representing AGB and BGB
 0.4682 = carbon fraction value referenced from Rahman et al. [18].
 CO₂e = carbon sequestration (kg/stand)

3. RESULTS AND DISCUSSION

3.1 Survival Rate

The planting of *R. mucronata mangroves* by PHE ONWJ on the coast of Pantai Bahagia Village shows excellent growth performance. It is indicated by the survival rate (R), which ranges from 98.67-100%. In the 2020-2021 range, 300 seedlings were observed to be alive (R = 100%); in the 2021-2022 range there were 296 out of 300 seedlings observed to live optimally (R = 98.67%), while in the 2022-2023 range, there were 295 out of 296 (R = 99.66%) seedlings alive and growing well (Fig. 2).




Fig. 2. The survival rate of *R. mucronata* seedlings from the REMAJA PHE ONWJ program in Pantai Bahagia Village, Bekasi – Indonesia

The indication of the success of planting *R. mucronata* mangroves was also seen in the significant diameter growth during the evaluation period (2020-2023). In the initial planting in 2020, the average seedling diameter of *R. mucronata* was 0.68 ± 0.0087 cm ($P > 0.05$). The diameter increased by 0.43 cm in 2021 (D2021 = 1.11 cm), 1.2 cm in 2022 (D2022 = 2.31 cm), and 0.45 cm in 2023 (D2023 = 2.76 cm) (Fig. 3). The growth rate between seedlings was relatively the same and significant at a confidence level of 95% ($\alpha = 0.05$).

The growth rate of *R. mucronata* planted by PHE ONWJ is greater than the natural growth on the West Muna coast, which is 0.6228 cm per year [15]. The growth rate of mangrove seedlings can be affected by several factors such as hydrological conditions and substrates. In addition, according to Armiani (2024), the presence of heavy metals such as lead (Pb) can inhibit the growth of *Rhizophora* sp seedlings, especially in diameter. The effect of heavy metals on the growth of mangrove seedlings has been previously reported by Ma and Yang [19] who found that the growth of *Candelia obovata* and *Bruguiera sexangula* is inhibited due to the accumulation of heavy metals. This is because the accumulation of heavy metals affects the physiology of mangroves, especially the ability of photosynthesis which ultimately inhibits the growth of mangroves [20,21].

The high survival rate and growth of *Rhizophora mucronata* mangroves on the coast of Pantai Bahagia Village is due to a combination of biological adaptation and supportive ecological conditions. *R. mucronata* can naturally adapt to extreme coastal environments, such as high salinity and anaerobic sediments, which allows them to grow optimally [11]. Good sediment quality and minimal biotic and abiotic disturbances also play an essential role in the success of the planting.

3.2 Mangrove Biomass

The average biomass value in 2020 (B2020) was 0.1554 kg/stand, consisting of 0.0825 kg AGB and 0.0729 kg BGB. This value continues to increase as the diameter grows every year. The biomass potential of *R. mucronata* based on its diameter growth is 0.4916 kg/stand (0.2752 kg AGB and 0.2164 kg BGB) in 2021, 2.8282 kg/stand (1.7058 kg AGB and 1.1224 kg BGB) in 2022, and 4.2321 kg/stand (2.5974 kg AGB and 1.6347 kg BGB) in 2023 (Fig. 4).

Based on the standing biomass, the total biomass potential contributed from the planting of 8000 seedlings of the REMAJA PHE ONWJ program is 1.24 tons in 2020 (N = 8000), 3.93 tons in 2021 (N = 8000), 22.33 tons in 2022 (N = 7894), and 33.29 tons in 2023 (N = 7867) (Table 1).




Fig. 3. Growth of *R. mucronata* seedling diameter ($P > 0.05$) from the REMAJA PHE ONWJ program on the coast of Pantai Bahagia Village, Bekasi – Indonesia

Table 1. Total biomass potential of *R. mucronata* from the REMAJA PHE ONWJ program on the coast of Pantai Bahagia Village, Bekasi – Indonesia

Years	N	AGB (ton)	BGB (ton)	Total (ton)
2020	8000	0.66	0.58	1.24
2021	8000	2.20	1.73	3.93
2022	7984	13.47	8.86	22.33
2023	7867	20.43	12.86	33.29

Notes: N is based on the survival rate, as shown in Figure 2; AGB is above-ground biomass, and BGB is below-ground biomass.




Fig. 4. The potential of standing biomass of *R. mucronata* from the REMAJA PHE ONWJ program in the Pantai Bahagia Village, Bekasi – Indonesia coast.

3.3 Carbon Stock and CO₂ Absorption

Based on the total biomass potential, planting *R. mucronata* mangroves from the REMAJA PHE ONWJ program significantly contributes to carbon storage. The results of the evaluation show that the total carbon stock stored in 2023 will reach 15.59 tons C, consisting of 9.57 tons of C on the above-ground (AGC) and 6.02 tons of C on the below-ground (BGC). The stock increased by 5.14 tons C compared to 2022, 13.75 tons C compared to 2021, and 15.01 tons C compared to 2020 (Fig. 5).

This significant increase in carbon stock reflects the success of the *Rhizophora mucronata* mangrove planting program as part of a climate change mitigation strategy. Mangroves are known to have an extraordinary ability to absorb and store carbon, both at the top of the soil (AGC) through stems, leaves, and branches and below the ground level (BGC) through roots [1,22]. The most significant contribution in 2023 came from AGC, which stores more carbon as

more mature and more extensive mangrove vegetation grows.

The role of mangroves in sequestering carbon provides short-term benefits and contributes to the stability of coastal ecosystems in the long term. Carbon stocks stored underground, such as those recorded in the BGC, can remain trapped for centuries if mangrove ecosystems are protected from damage. With an annual upward trend in carbon storage, programs like REMAJA PHE ONWJ not only support environmental conservation but also help reduce greenhouse gas emissions, making mangrove planting areas an integral part of nature-based solutions to climate change [23].

The contribution to reducing greenhouse gas emissions, especially CO₂, can be equivalent based on the potential of carbon stocks [22,24]. Based on this, the contribution of *R. mucronata* absorption in the 2020-2021, 2021-2022, and 2022-2023 ranges were 4.62 tons of CO₂e, 31.58 tons of CO₂e, and 18.83 tons of CO₂e, respectively (Fig. 6).




Fig. 5. Potential carbon stock of *R. mucronata* from the REMAJA PHE ONWJ program on the coast of Pantai Bahagia Village, Bekasi – Indonesia.




Fig. 6. Carbon sequestration rate (CO₂e) of *R. mucronata* from the REMAJA PROGRAM PHE ONWJ program on the coast of Pantai Bahagia Village, Bekasi – Indonesia

4. CONCLUSION

The planting of *R. mucronata* through the REMAJA PHE ONWJ program in the Karawang Regency's coastal area significantly contributes to increasing carbon stocks. From 2020 to 2023, the total carbon stock stored increased consistently, reaching 15.59 tonnes C in 2023, 9.57 tonnes C at above-ground carbon (AGC), and 6.02 tonnes C at below-ground carbon (BGC). This increase reflects the program's success in supporting climate change mitigation through long-term carbon storage, with carbon stocks continuing to grow as more mature mangrove vegetation grows.

The program also demonstrated its effectiveness in maintaining a high survival rate of mangrove seedlings, with a survival rate of more than 98% throughout the evaluation period. In addition, the increase in mangrove biomass, both above and below ground, contributes to a more significant total carbon stock year over year. Overall, the planting of *R. mucronata* under this program provides ecological benefits in the form of carbon storage and supports the long-term stability of coastal ecosystems while serving as a natural solution to reduce greenhouse gas emissions.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image

generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that they have no known competing financial interests or non-financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

1. Rahman, Ceantury A, Tuahatu JW, Lokollo FF, Supusepa J, Hulopi M, Permatahati YI, Lewerissa A, Wardiatno Y. Mangrove ecosystem in Southeast Asia region: mangrove extent, blue carbon potential and CO₂ emission in 1996-2020. *Science of the Total Environment*. 2024a;915(3):1-12.
2. Alongi DM. Carbon cycling and storage in mangrove forests. *Annual Review of Marine Science*. 2014;195-219. Available:DOI: 10.1146/annurev-marine-010213-135020
3. Murdiyarsa D, Purbopuspito J, Kauffman J. The potential of Indonesian mangrove forests for global climate change mitigation. *Nat. Clim. Change*. 2015;5: 1089-1092. Available:https://doi.org/10.1038/nclimate2734

4. Donato DC, Kauffman JB, Murdiyarso D, Kurnianto S, Stidham M, Kannien M. Mangroves among the most carbon-rich forests in the tropics. *Nature Geosci.* 2011;4:293–297.
Available:<https://doi.org/10.1038/ngeo1123>
5. Adame MF, Santini NS, Tovilla C, Lule AV, Castro L, Guevara M. Carbon stock and soil sequestration rates of tropical riverine wetlands. *Biogeosci.* 2015;12:3805-3818.
Available:<https://doi.org/10.5194/bg-12-3805-2015>
6. Sidik F, Lawrence A, Wagey T, Zamzani F, Lovelock CE. Blue carbon: A new paradigm of mangrove conservation and management in Indonesia. *Marine Policy.* 2023;147: 1-9.
7. Ilman M, Dargusch P, Dart P, Onrizal. A historical analysis of the drivers of loss and degradation Indonesia's mangroves. *Land Use Policy.* 2016;54:448-459.
Available:<https://doi.org/10.1016/j.landusepol.2016.03.010>
8. Rahman, Wardiatno Y, Yulianda F, Rusmana I, Bengen DG. *Metode dan Analisis Studi Ekosistem Mangrove*. Bogor (ID): IPB Press. 2020;124.
9. Amalo LF, Setiawan Y, Kusmana C. Spatio-temporal dynamics of mangrove extent in coastal Bekasi Regency, Indonesia. *Proceeding IOP Conference Series: Earth and Environmental Science.* 2023;1266(012005):1-9.
10. Djamaluddin R. *Mangrove: Biologi, Ekologi, Rehabilitasi, dan Konservasi*. Unsrat Press. 2018;237.
11. Bengen DG, Yonvitner Rahman. *Pedoman Teknis Pengenalan dan Pengelolaan Ekosistem Mangrove*. IPB Press. 2022 ;88.
12. Samal, P, Srivastava J, Charles B., Singarasubramaniam SR. Species distribution models to predict the potential niche shift and priority conservation areas for mangroves (*Rhizophora apiculata*, *R. mucronata*) in response to climate and sea level rise fluctuations along coastal India. *Ecological Indicators.* 2023;154:1-16.
13. Rahman, Wardiatno Y, Yulianda F, Rusmana I. Production ratio of seedlings and density status of mangrove ecosystems in coastal areas of Indonesia. *Advances in Environmental Biology.* 2019;13(6):13-21.
14. Rahman, Lokollo FF, Manuputty GD, Hukubun RD, Krisye, Maryono, Wawo M, Wardiatno Y. A review on the biodiversity and conservation of mangrove ecosystems in Indonesia. *Biodiversity and Conservation.* 2024b;33(3):875-903.
15. Rahman. *Model pengelolaan ekosistem mangrove berbasis dinamika stok karbon dan fluks gas rumah kaca*. Bogor. [Disertation]. IPB University. 2020;110.
16. Komiyama A, Poungporn S, Kato S. Common allometric equation for estimating the tree weight of mangroves. *J. Trop. Ecol.* 2005;21:471–477.
Available:<https://doi.org/10.1017/S0266467405002476>
17. World Agroforestry Center. wood density of *R. mucronata*; 2024.
Available:http://db.worldagroforestry.org/wd/species/Rhizophora_mucronata. In September 2024.
18. Rahman, Maryono, Sigiyo ON. What is the true carbon fraction of mangrove biomass? *Malaysian Journal of Science.* 2023;42(2): 1 – 6.
Available:<https://doi.org/10.22452/mjs.vol42no2.10>
19. Ma Li, Yang S. Growth and physiological response of *Kandelia obovata* and *Bruguiera sexangula* seedlings to aluminum stress. *Environ. Sci. Pollut. Res. Int.* 2022;29(28):43251–43266.
Available:<https://doi.org/10.1007%2Fs11356-021-17926-0>
20. MacFarlane GR, Burchett MD. Toxicity, growth and accumulation relationships of copper, lead and zinc in the grey mangrove *Avicennia marina* (Forsk.) Vierh. *Mar. Environ. Res.* 2002;54(1):65 – 84.
Available:[https://doi.org/10.1016/s0141-1136\(02\)00095-8](https://doi.org/10.1016/s0141-1136(02)00095-8)
21. Parida AK, Jha B. Salt tolerance mechanisms in mangroves: a review. *Trees.* 2010;24:199–217.
Available:<https://doi.org/10.1007/s00468-010-0417-x>
22. Rahman, Tuahatu JW, Tuhehay. Blue carbon potential on mangrove stands on the coast of Waai Village, Central Maluku Regency. *J. Ilmu Kehutanan.* 2024c;18(1):7-15.

23. Huxham M, Kairu A, Lang'at JA, Kivugo R, Mwafrica M, Huff A, Shilland R. Rawls in the mangrove: Perceptions of justice in nature-based solutions projects. *People and Nature*. 2023;1-16. Available: <https://doi.org/10.1002/pan3.10498>
24. Armiani S. Pengaruh pemberian berbagai konsentrasi timbal (Pb) terhadap pertumbuhan bibit mangrove *Rhizophora stylosa*. *Panthera*. 2024;4(3):112 – 120.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/124263>

Full Article - PDF

Review History

Discussion

Published: 2024-10-03

DOI: 10.9734/ajee/2024/v23i10615

Page: 121-129

Issue: 2024 - Volume 23 [Issue 10]



? Total citations
? Recent citations
n/a Field Citation Ratio
n/a Relative Citation Ratio

?

Evaluating Carbon Stocks (2020-2023) in *Rhizophora mucronata* Plantations under the PHE ONWJ REMAJA Program in Pantai Bahagia Villages, Bekasi Regency, Indonesia

Rahman

Department of Marine Science, Faculty of Fisheries and Marine Science, Pattimura University, Indonesia.

Luisa Febrina Amalo *

The Central for Environment Research, IPB University, West Java, Bogor, Indonesia.

Pungki Ari Wibowo

The Central for Environment Research, IPB University, West Java, Bogor, Indonesia.

Luluk Dwi Wulan Handayani

The Central for Environment Research, IPB University, West Java, Bogor, Indonesia.

Hadi Supardi

Department of Environmental, HSSE Division, PT. Pertamina Hulu Energi Offshore North West Java, Jakarta, Indonesia.

L. M. Alfin Agushara Bena

Department of Environmental, HSSE Division, PT. Pertamina Hulu Energi Offshore North West Java, Jakarta, Indonesia.

Arif Rahman

Department of Environmental, HSSE Division, PT. Pertamina Hulu Energi Offshore North West Java, Jakarta, Indonesia.

*Author to whom correspondence should be addressed.

Abstract

This study evaluates the carbon storage potential of *Rhizophora mucronata* plantations as part of the North Java Coast Mangrove Restoration Program (REMAJA) implemented by PT. Pertamina Hulu Energi Offshore North West Java (PHE ONWJ) in Pantai Bahagia Village, Bekasi Regency. A total of 8,000 mangrove seedlings were planted in 2020, and the diameter growth and carbon stock were monitored from 2020 to 2023. The monitoring process involved measuring the trunk diameter of the mangrove, combined with allometric equations to estimate biomass. Carbon sequestration was calculated by applying conversion factors to the biomass values, allowing the quantification of stored carbon. This method provides an accurate assessment of the mangrove's role in carbon sequestration over time. The results showed that the survival rate of *Rhizophora mucronata* seedlings was exceptionally high, averaging over 98%. The total carbon stock increased annually, reaching 15.59 tons of carbon (C) in 2023, 9.57 tons in AGC, and 6.02 tons in BGC. The result represents a significant increase compared to previous years, with an additional 5.14 tons of carbon compared to 2022, 13.75 tons more than in 2021, and 15.01 tons more than in 2020. Based on these results, the carbon dioxide sequestration contribution of *R. mucronata* during the periods of 2020-2021, 2021-2022, and 2022-2023 was 4.62 tons CO₂e, 31.58 tons CO₂e, and 18.83 tons CO₂e, respectively. In conclusion, the *R. mucronata* plantation under this program significantly enhanced carbon stocks, supporting climate change mitigation through long-term carbon storage. This program demonstrates that mangrove rehabilitation can play a crucial role in protecting coastal ecosystems and reducing greenhouse gas emissions.

Keywords: Carbon stock, climate change mitigation, *R. mucronata*, mangrove rehabilitation

HOW TO CITE

Rahman, Luisa Febrina Amalo, Pungki Ari Wibowo, Luluk Dwi Wulan Handayani, Hadi Supardi, L. M. Alfin Agushara Bena, and Arif Rahman. 2024. "Evaluating Carbon Stocks (2020-2023) in *Rhizophora Mucronata* Plantations under the PHE ONWJ REMAJA Program in Pantai Bahagia Villages, Bekasi Regency, Indonesia". *Asian Journal of Environment & Ecology* 23 (10):121-29. <https://doi.org/10.9734/ajee/2024/v23i10615>.

More Citation Formats ▾

[Home](#) / [About the Journal](#)

About the Journal

ISSN: 2456-690X

Journal DOI: <http://dx.doi.org/10.9734/ajee> [\(Link\)](#)

Peer-review model: [Advanced OPEN peer review](#)

Digital Archiving: [Journal Repository \(JR\)](#), [LOCKSS](#), [CLOCKSS](#), [KEEPERS](#)

NAAS Score: 4.76 (2024) (Proof: <https://naas.org.in/NJS/journals2024.pdf>)

Asian Journal of Environment & Ecology (ISSN: 2456-690X) aims to publish high quality papers ([Click here for Types of paper](#)) in all areas of 'Environment and Ecology'. By not excluding papers based on novelty, this journal facilitates the research and wishes to publish papers as long as they are technically correct and scientifically motivated. The journal also encourages the submission of useful reports of negative results. This is a quality controlled, OPEN peer-reviewed, open-access INTERNATIONAL journal.

This is an open-access journal which means that all content is freely available without charge to the user or his/her institution. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author. This is in accordance with the BOAI definition of open access.

Every issue will consist of a minimum of 5 papers. Each issue will be running, and all officially accepted manuscripts will be immediately published online. The state-of-the-art running issue concept gives authors the benefit of 'Zero Waiting Time' for the officially accepted manuscripts to be published. This journal is an international journal and its scope is not confined by the boundary of any country or region.

Disclaimer: This international journal has no connection with any scholarly society or association or any specific geographic location or any country (like USA, UK, Germany, etc). This is an independent journal. Legal contact address of editorial office and other offices are available in this [link](#).